

Key Kid Starter 18% Pellet R

9303253

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christenseni*, and *E. ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin Sodium 30 g/ton

Complete Feed for Goats

INGREDIENTS

Processed Grain By-Products, Grain Products, Plant Protein Products, Roughage Products, Forage Products, Calcium Carbonate, Molasses Products, Salt, Ammonium Chloride, Maxi Bond, Potassium Magnesium Sulfate, Propionic acid, Magnesium Oxide, Dicalcium Phosphate, Sodium Selenite, Mineral Oil, Manganous Hydroxychloride, Zinc Hydroxide, Vitamin E Supplement, Basic Copper Chloride, Vitamin A Acetate, Vit D3 Supplement, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Natural Flavors., Zinc Sulfate, Manganous Oxide, Ferrous Sulfate, Vitamin D3 Supplement, Vitamin B12 Supplement, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Yeast Culture, Yeast Extract, Hydrolyzed Yeast.

FEEDING DIRECTIONS

Feed Key Kid Starter 18% Pellet R with good quality forages and/or roughages to meet the nutritional requirements of growing goats. Free choice salt may be provided. Supply plenty of fresh clean water at all times. For a more specific feeding program, please contact your Keystone Mills Feed Consultant.

CAUTION: Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet.

CAUTION: USE AS DIRECTED

THIS FEED SHOULD BE USED ONLY IN ACCORDANCE WITH THE RECOMMENDATIONS ON THIS FEED TAG

CAUTION: Do not allow horses or other equines access to feeds containing Monensin.

Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats.

Monensin medicated feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions.

LIMITATIONS: Feed continuously. Do not feed to lactating goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the level of Monensin recommended in the feeding direction, as reduced average daily gains may result.

FDA REGULATION FOR MONENSIN

Re-circulation or Agitation required before use: Re-circulating tank systems: Re-circulate immediately prior to use for no less than 10 minutes, moving not less than 1 percent of the tank contents per minute from the bottom of the tank to the top. Re-circulate daily as described even when not used. Agitation tank systems (mechanical, air, etc.): Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily as described even when not used.

EXPIRATION DATE MARKED ON TAG: (Must be used within 8 weeks of manufacture)

WARNING: Must Observe feeding directions on label.

MANUFACTURED BY:



KEYSTONE MILLS

— Serving Agribusiness —

Romulus, NY 14541

(315) 549-8226

NET WT. 50 lbs (22.68 kg)

Lot#:01-000020219658-324

rev: 09/29/17