

# Key Dry Cow 26% Pellet R

9706603

## MEDICATED

For improved feed efficiency.

ACTIVE DRUG INGREDIENT  
Monensin Sodium .... 117 g/ton

Feed for Non-Lactating Dairy Cattle

### GUARANTEED ANALYSIS

Crude Protein	(Min)	27.00 %
---------------	-------	---------

\* (This Includes not more than 1.4 % equivalent non-protein nitrogen.)

Crude Fat	(Min)	4.10 %
Crude Fiber	(Max)	6.50 %
Calcium (Ca)	(Min) 0.48 (Max)	1.18 %
Phosphorus (P)	(Min)	0.80 %

### INGREDIENTS

Processed Grain By-Products, Plant Protein Products, Grain Products, Molasses Products, Magnesium Sulfate, Magnesium Oxide, Calcium sulfate, Monosodium Phosphate XP4 (50#), Mono-Dicalcium Phosphate, Urea, Vitamin E Supplement, Calcium Carbonate, Maxi Bond, Selenium yeast, Brewers dried yeast., Yeast Culture, Yeast Extract, Hydrolyzed Yeast, Manganous Hydroxychloride, Zinc Hyroxchloride, Basic Copper Chloride, Vitamin A Acetate, Vit D3 Supplement, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Natural Flavors..

### FEEDING DIRECTIONS

Feed Key Dry Cow 26% Pellet R with good quality forages and/or roughages to meet the nutritional requirements of dry dairy cattle. Provide plenty of fresh clean water at all times. For a more specific feeding program, please contact your Keystone Mills Dairy Consultant.

CAUTION: Do not feed supplemental selenium in excess of 0.3 ppm in the total nutritional diet.

CAUTION: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result.

Limitations: For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

WARNING: Must Observe feeding directions on label.

MANUFACTURED BY:



**KEYSTONE MILLS**

*Serving Agriculture in America*

Romulus, NY 14541

(315) 549-8226

NET WT. 50 lbs (22.68 kg)

rev: 05/20/13

Lot# 01-0000222553-12